

Squeezed primordial bispectrum from general vacuum state

Jinn-Ouk Gong^{a,b,*} and Misao Sasaki^{c,†}

^a*Asia Pacific Center for Theoretical Physics, Pohang 790-784, Korea*

^b*Department of Physics, Postech, Pohang 790-784, Korea*

^c*Yukawa Institute for Theoretical Physics,
Kyoto University, Kyoto 606-8502, Japan*

(Dated: April 2, 2013)

We study the general relation between the power spectrum and the squeezed limit of the bispectrum of the comoving curvature perturbation produced during single-field slow-roll inflation when the initial state is a general vacuum. Assuming the scale invariance of the power spectrum, we derive a formula for the squeezed limit of the bispectrum, represented by the parameter f_{NL} , which is not slow-roll suppressed and is found to contain a single free parameter for a given amplitude of the power spectrum. Then we derive the conditions for achieving a scale-invariant f_{NL} , and discuss a few examples.

I. INTRODUCTION

Currently inflation is regarded as the leading candidate to provide the initial conditions for the hot big bang evolution of the universe [1]. During inflation the primordial curvature perturbation is generated, which after inflation becomes seeds for the temperature fluctuations in the cosmic microwave background (CMB) and the large scale structures of the universe. Recent observations [2] are consistent with the predictions of inflation, i.e. the primordial fluctuations are statistically almost perfectly Gaussian with a nearly scale invariant power spectrum. Thus, one of the main tasks of ongoing and future observation programmes such as PLANCK [3] is to test if there is any deviation from these predictions. These high precision future observations will able us to rule out and/or further constrain various models of inflation, thus shedding light on the physics of the very early universe.

Among observable signatures, non-Gaussianity has been attracting great interest. In particular a lot of efforts have been made to detect a non-zero three-point correlation function, or its Fourier transform, the bispectrum, of the primordial perturbation [4]. The bispectrum is specified by three parameters, and templates for various configurations in the momentum space have been proposed

* jinn-ouk.gong@apctp.org

† misao@yukawa.kyoto-u.ac.jp

and used in observation. Among them, a particularly useful one is the squeezed configuration, where one of three momenta is much smaller than the others, e.g. $k_1 \approx k_2 \gg k_3$. A dominant source of non-Gaussianity for this configuration is the so-called local one, where the curvature perturbation is locally expanded as [5, 6]

$$\mathcal{R}(x) = \mathcal{R}_g(x) + \frac{3}{5}f_{\text{NL}}\mathcal{R}_g^2(x) + \cdots, \quad (1)$$

where the subscript g denotes the dominant Gaussian component. The coefficient f_{NL} determines the size of non-Gaussianity in the bispectrum.

An important prediction of single-field slow-roll inflation is that, in the squeezed limit, f_{NL} is proportional to the spectral index $n_{\mathcal{R}} - 1$ of the power spectrum [6, 7], and is thus too small to be observed. This relation holds irrespective of the detail of models and is usually called the consistency relation. Thus, barring the possibility of features that correlate the power spectrum and f_{NL} [8], it has been widely claimed that any detection of the local non-Gaussianity would rule out *all* single field inflation models. However, it is based on two assumptions. First, the curvature perturbation is frozen outside the horizon and does not evolve. That is, only one growing mode is relevant on super-horizon scales. Indeed, it is possible to make use of the constancy of $\mathcal{R}$ to extract only a few relevant terms in the cubic order Lagrangian to simplify considerably the calculation of the squeezed bispectrum, and to confirm the consistency relation [9]. If we abandon this assumption, the usual consistency relation does not hold any longer [10].

Another assumption is that deep inside the horizon interactions are negligible and the state approaches the standard Fock vacuum in the Minkowski space, so-called the Bunch-Davies (BD) vacuum. If this assumption does not hold, the corresponding bispectrum may be enhanced in the folded limit [11], in particular in the squeezed limit [12, 13]. Thus, the usual consistency relation may not hold. See also [14], where the violation of the tree level consistency relation is discussed together with the infrared divergence in the power spectrum from one-loop contributions for non-BD initial states. However, in the previous studies the relation between the power spectrum and bispectrum was unclear and f_{NL} was not easily readable [12], or case studies on specific models were carried out [13]. It is then of interest to make a closer and more explicit study on the general relation between the power spectrum and the squeezed limit of the bispectrum.

In this article, we compute the squeezed limit of the bispectrum when the initial state is not the BD vacuum and study the relation between the squeezed limit of the primordial bispectrum, described by the non-linear parameter f_{NL} and the power spectrum. We find indeed f_{NL} can be significantly large, but its momentum dependence is in general non-trivial. We then discuss the

condition for f_{NL} to be momentum-independent, thus exactly mimics the local form (1).

Before proceeding to our analysis, let us make a couple of comments. First, we note that the squeezed limit does not necessarily mean the exact limit of a squeezed triangle in the momentum space. It includes the case when the wavenumber of the squeezed edge of a triangle is smaller than that of the observationally smallest possible wavenumber, i.e. that corresponds to the current Hubble parameter. In the context of (1), it needs to be valid only over the region covering our current Hubble horizon size. Second, in our analysis we focus only on the squeezed limit of the bispectrum and its relation to the power spectrum. However, if a large f_{NL} that mimics the local form of the non-Gaussianity is generated, we may also have the bispectrum with a non-negligible amplitude in some other shapes of the triangle [12, 13]. This may be an interesting issue to be studied, but it is out of the scope of this work.

II. BISPECTRUM IN SINGLE-FIELD SLOW-ROLL INFLATION

For general single-field inflation, the equation of motion of the comoving curvature perturbation $\mathcal{R}$ is given by [15]

$$(z^2 \mathcal{R}'_k)' + c_s^2 k^2 z^2 \mathcal{R}_k = 0, \quad (2)$$

where a prime denotes a derivative with respect to the conformal time $d\tau = dt/a$, $\epsilon \equiv -\dot{H}/H^2$ and $z^2 \equiv 2m_{\text{Pl}}^2 a^2 \epsilon / c_s^2$ and c_s is the speed of sound. From (2), we can see that irrespective of the detail of the matter sector, a constant solution of $\mathcal{R}_k$ always exists on super-sound-horizon scales, $c_s k \ll aH$, and it dominates at late times for slow-roll inflation for which $z^{-1} \sim a^{-1} \sim \tau$.

Here we focus on the case of slow-roll inflation. Keeping the constancy of $\mathcal{R}_k$ on large scales, in the squeezed limit $k_1 \approx k_2$ and $k_3 \rightarrow 0$, the bispectrum at $\tau = \bar{\tau}$ is given by [9]

$$B_{\mathcal{R}}(k_1, k_2, k_3; \bar{\tau}) = \left[\frac{\eta(\bar{\tau})}{c_s^2(\bar{\tau})} + \frac{F(k_1, \bar{\tau})}{P_{\mathcal{R}}(k_1)} \right] P_{\mathcal{R}}(k_1) P_{\mathcal{R}}(k_3), \quad (3)$$

$$F(k, \bar{\tau}) = i\mathcal{R}_k^2(\bar{\tau}) \int_{-\infty}^{\bar{\tau}} d\tau \left[\frac{2\epsilon}{c_s^4} (\epsilon - 3 + 3c_s^2) a^2 (\mathcal{R}_k'^*)^2 + \frac{2\epsilon}{c_s^2} (\epsilon - 2s + 1 - c_s^2) a^2 k^2 (\mathcal{R}_k^*)^2 \right. \\ \left. + \frac{2\epsilon}{c_s^2} \frac{d}{d\tau} \left(\frac{\eta}{c_s^2} \right) a^2 \mathcal{R}_k'^* \mathcal{R}_k^* + \text{c.c.} \right], \quad (4)$$

where $\eta \equiv \dot{\epsilon}/(H\epsilon)$ and $s \equiv \dot{c}_s/(Hc_s)$.

Being interested in large non-Gaussianity, among the terms inside the square brackets of (4) we may focus on those not suppressed by the slow-roll parameters, that is,

$$F_0 = \frac{2(1 - c_s^2)}{c_s^2} \Re \left\{ i\mathcal{R}_k^2(\bar{\tau}) \int_{-\infty}^{\bar{\tau}} d\tau \left[-3z^2 (\mathcal{R}_k'^*)^2 + c_s^2 k^2 z^2 (\mathcal{R}_k^*)^2 \right] \right\}, \quad (5)$$

where for simplicity we have assumed the time variation of c_s^2 is negligible, $s = 0$. Now, we find it is more convenient to write the integrand of (5) in terms of $\mathcal{R}'_k$. Multiplying (2) by $\mathcal{R}_k$, we have

$$c_s^2 k^2 z^2 \mathcal{R}_k^2 = - (z^2 \mathcal{R}'_k \mathcal{R}_k)' + (z \mathcal{R}'_k)^2 . \quad (6)$$

Hence (5) becomes

$$F_0 = \frac{2(1-c_s^2)}{c_s^2} \Re \left\{ i \mathcal{R}_k^2(\bar{\tau}) \left[-z^2 \mathcal{R}_k^* \mathcal{R}_k^{*'}(\bar{\tau}) - 2 \int_{-\infty}^{\bar{\tau}} d\tau (z \mathcal{R}_k^{*'})^2 \right] \right\} \equiv \frac{2(1-c_s^2)}{c_s^2} (\Re[I_1] + \Re[I_2]) . \quad (7)$$

As we can write $\mathcal{R}_k$ in terms of $\mathcal{R}'_k$ as (2), we do not have to work with $\mathcal{R}_k$ but only need to solve for $\mathcal{R}'_k$. Setting

$$f \equiv z \mathcal{R}'_k , \quad (8)$$

and taking a derivative of (2), we obtain

$$f'' + [c_s^2 k^2 - z(z^{-1})''] f = 0 . \quad (9)$$

An interesting property of this equation is that in the slow-roll case, $z^{-1} \sim a^{-1} \sim \tau$, so the potential term $z(z^{-1})''$ vanishes at leading order [16]. Specifically we have

$$z(z^{-1})'' = a^2 H^2 \left[\epsilon + \frac{\eta}{2} + \mathcal{O}(\eta^2, \epsilon\eta) \right] . \quad (10)$$

This means that the WKB solution $f \propto e^{\pm i c_s k \tau}$ remains valid even on super-sound-horizon scales at leading order in the slow-roll expansion. The general leading order solution during slow-roll inflation is thus

$$f = \sqrt{\frac{c_s k}{2}} \left(C_k e^{-i c_s k \tau} + D_k e^{i c_s k \tau} \right) , \quad (11)$$

where C_k and D_k are constant and we have extracted the factor $\sqrt{c_s k/2}$ for convenience.

Carrying out the standard quantization procedure, we find that for f to be properly normalized, the constants C_k and D_k satisfy

$$|C_k|^2 - |D_k|^2 = 1 . \quad (12)$$

Setting $D_k = 0$ corresponds to the usual choice of the BD vacuum. But here we do not assume so and let D_k be generally non-zero. From (6) and (11), the power spectrum can be easily computed to be

$$\mathcal{P}_{\mathcal{R}} = \frac{k^3}{2\pi^2} |\mathcal{R}_k|_{c_s k = aH}^2 = \frac{k^3}{2\pi^2} \frac{1}{2(c_s k)^3} \left(\frac{z'}{z^2} \right)_{c_s k = aH}^2 |C_k + D_k|^2 . \quad (13)$$

Thus, a scale-invariant spectrum requires $|C_k + D_k|^2 \sim k^0$.

Now we return to (7). For slow-roll inflation, $\mathcal{R}'_k$ rapidly decays outside the sound horizon. However, since z grows like a , neither I_1 nor I_2 may not be negligible outside the sound horizon. Rewrite them in terms of f , we easily find the expression for $\Re[I_1]$ as

$$\Re[I_1] = \Re \left[\frac{i}{z^2 (c_s k)^6} \left| f' + \frac{z'}{z} f \right|^2 f' f^* \right] = \frac{k^{-3}}{c_s^5} \left(\frac{z'}{z^2} \right)^2 |C_k + D_k|^2, \quad (14)$$

where we have used (12). The second term is

$$\begin{aligned} \Re[I_2] &= \Re \left[-i \frac{2}{z^2 (c_s k)^4} \left(f' + \frac{z'}{z} f \right)^2 \int_{-\infty}^{\bar{\tau}} d\tau (f^*)^2 \right] \\ &\approx -\frac{k^{-4}}{z^2 c_s^5} \Re \left[\left(f' + \frac{z'}{z} f \right)^2 \left(C_k^{*2} e^{2ic_s k \bar{\tau}} + D_k^{*2} e^{-2ic_s k \bar{\tau}} - 4ic_s k \tau_\infty C_k^* D_k^* \right) \right]. \end{aligned} \quad (15)$$

Here, upon integrating the last term of the integrand, there is no time dependence and thus literally integrating from $-\infty$ it diverges. However in reality it should be understood as the boundary τ_∞ with $|\bar{\tau}| \ll |\tau_\infty|$ at which the initial condition is specified. This means depending on our choice of $c_s k \tau_\infty$, the contribution of this term may become very large, in fact can be made arbitrarily large. Hence we cannot neglect it even in the limit $k \rightarrow 0$. In this limit,

$$\Re[I_2] = -\frac{k^{-3}}{c_s^5} \left(\frac{z'}{z^2} \right)^2 \Re \left[(C_k + D_k)^2 \left(C_k^{*2} - D_k^{*2} - 4ic_s k \tau_\infty C_k^* D_k^* \right) \right]. \quad (16)$$

If we only consider the contribution from the terms $C_k^{*2} - D_k^{*2}$, the result is precisely $-\Re[I_1]$ and hence cancels out. For the other terms in (4), the calculation goes more or less the same, and we find slow-roll suppressed contributions are given in the form, $\Re[I_1] + \Re[I_2] = \epsilon P_{\mathcal{R}}(k) \left[1 - 2c_s k \tau_\infty \Re(i C_k D_k^*) |C_k + D_k|^{-2} \right]$.

Thus, from (3) in the squeezed limit the additional contribution to the non-linear parameter f_{NL} when $D_k \neq 0$ is given by

$$\frac{3}{5} f_{\text{NL}} = \frac{F_0}{4P_{\mathcal{R}}(k)} = \left(\frac{1 - c_s^2}{c_s^2} - \frac{\epsilon}{c_s^2} \right) c_s k \tau_\infty \frac{\Re(i C_k D_k^*)}{|C_k + D_k|^2}. \quad (17)$$

Note that the only assumption we have made is slow-roll inflation where $z^{-1} \sim a^{-1} \sim \tau$, and thus all the above arguments are completely valid for general vacuum state under the constancy of the curvature perturbation $\mathcal{R}$.

III. LOCAL, SCALE-INDEPENDENT f_{NL}

From (17), we see that f_{NL} will be k -dependent in general due to that of $C_k D_k^*$, in addition to that from non-linear evolution on large scales [17]. With the normalization (12), we may

parametrize C_k and D_k as

$$C_k = e^{i\alpha_k} \cosh \chi_k, \quad (18)$$

$$D_k = e^{i\beta_k} \sinh \chi_k, \quad (19)$$

From the power spectrum (13), by setting $\mathcal{A} \equiv |C_k + D_k|^2$ which should be almost k -independent, we can solve for χ_k as

$$\sinh(2\chi_k) = \frac{\mathcal{A}^2 + \mathcal{A}\sqrt{\mathcal{A}^2 - \sin^2 \varphi_k} - \cos \varphi_k - 1}{(1 + \cos \varphi_k) \left(\mathcal{A} + \sqrt{\mathcal{A}^2 - \sin^2 \varphi_k} \right)}, \quad (20)$$

where $\varphi_k = \alpha_k - \beta_k$.

Meanwhile, for f_{NL} (17) we have, extracting the only (possibly) scale dependent part,

$$\mathcal{B} \equiv -c_s k \tau_\infty \Re(iC_k D_k^*) = \frac{1}{2} c_s k \tau_\infty \sin \varphi_k \sinh(2\chi_k). \quad (21)$$

With a suitable cutoff τ_∞ , we may choose φ_k and χ_k to make (21) have a particular k -dependence. Further, given the amplitude of the power spectrum $\mathcal{A}$, $\sin(2\chi_k)$ is written in terms of φ_k as (20) so f_{NL} contains a single free parameter φ_k other than the cutoff. To proceed further, let us for illustration consider two different choices of τ_∞ , and see when f_{NL} becomes scale-invariant. These choices are depicted in Figure 1.

Note that the conditions we derive below are phenomenological ones to be satisfied if f_{NL} is to remain almost scale-invariant. One may well try to construct more concrete and realistic models which can be approximated to the cases below, but the construction of such models is beyond the scope of the present paper.

(A) $\tau_\infty = -1/(c_s k_\infty)$:

This corresponds to fixing τ_∞ common to all modes. This will be the case when there is a phase transition at $\tau = \tau_\infty$ [18]. In this case, $-c_s k \tau_\infty = k/k_\infty \gg 1$ and we can think of three simple possibilities that give k -independent f_{NL} :

1. $\varphi_k \ll 1$: In this case we find

$$\mathcal{B} \approx -\frac{1}{2} \frac{k}{k_\infty} \varphi_k \sinh(2\chi_k) \approx -\frac{\mathcal{A}^2 - 1}{4\mathcal{A}} \frac{k}{k_\infty} \varphi_k. \quad (22)$$

Thus, by choosing $\varphi_k = \gamma_c k_\infty/k$, with γ_c being constant, we can make f_{NL} scale-invariant.

2. $2\chi_k \ll 1$: Likewise, we find

$$\mathcal{B} \approx -\frac{1}{2} \frac{k}{k_\infty} 2\chi_k \sin \varphi_k. \quad (23)$$




FIG. 1: Schematic plot of the two choices of the cutoff τ_∞ . On the left panel it is fixed as a constant $\tau_\infty = -1/(c_s k_\infty)$ and is the same for every mode. Meanwhile on the right panel it varies for different k in such a way that it corresponds to a fixed length scale, $c_s k |\tau_\infty| = \gamma_p$.

Thus $2\chi_k \approx \gamma_p k_\infty/k$ with a k -independent φ_k works as well. Note that in this case, φ_k is constant but its value is not constrained, and $\mathcal{A} \approx 1$ so that the state is very close to the BD vacuum.

3. $\varphi_k \ll 1$ and $2\chi_k \ll 1$: We have

$$\mathcal{B} \approx -\frac{1}{2} \frac{k}{k_\infty} \varphi_k 2\chi_k. \quad (24)$$

Thus, choosing $\varphi = p(k_\infty/k)^n$ and $2\chi_k = q(k_\infty/k)^{1-n}$ with p, q and $0 < n < 1$ being constant gives $\mathcal{B} = -pq/2$, so that f_{NL} is k -independent.

(B) $\tau_\infty = -\gamma_p/(c_s k)$ with $\gamma_p \gg 1$:

In this case, the cutoff τ_∞ depends on k in such a way that $-c_s k \tau_\infty = \gamma_p$ is constant. This is the case when the cutoff corresponds to a fixed, very short physical distance. Hence this cutoff may be relevant when we consider possible trans-Planckian effects [19]. Again, let us consider three simple possibilities:

1. $\varphi_k \ll 1$: We obtain

$$\mathcal{B} \approx -\frac{\gamma_p}{2} \varphi_k \sinh(2\chi_k) \approx -\frac{\gamma_p (\mathcal{A}^2 - 1)}{4\mathcal{A}} \varphi_k. \quad (25)$$

Thus we require φ_k to have no k -dependence in order to have a scale-invariant f_{NL} .

2. $2\chi_k \ll 1$: In this case $\mathcal{B} \approx -(\gamma_p/2)2\chi_k \sin \varphi_k$. Thus it is k -independent if both φ_k and χ_k are constant, for an arbitrary value of φ_k .
3. $\varphi_k \ll 1$ and $2\chi_k \ll 1$: This gives

$$\mathcal{B} \approx -\frac{\gamma_p}{2}\varphi_k 2\chi_k. \quad (26)$$

This is a limiting case of the second case above, and the simplest example is when both φ_k and $2\chi_k$ are k -independent.

We note that in all the cases considered above, f_{NL} can be large, say $f_{\text{NL}} \gtrsim 10$, if $c_s^2 \neq 1$ and the constant γ_c or γ_p is large.

IV. CONCLUSION

In this article, focusing on single-field slow-roll inflation, we have studied in detail the squeezed limit of the bispectrum when the initial state is a general vacuum. In this case, the standard consistency relation between the spectral index of the power spectrum of the curvature perturbation and the amplitude of the squeezed limit of the bispectrum does not hold. In particular, the squeezed limit of the bispectrum may not be slow-roll suppressed.

Under the assumption that the comoving curvature perturbation is conserved on super-sound-horizon scales, we have derived the general relation between the squeezed limit of the primordial bispectrum, described in terms of the non-linear parameter f_{NL} and the power spectrum. We find f_{NL} is indeed not slow-roll suppressed. But it depends explicitly on the momentum in general, hence may not be in the local form. We then have discussed the condition for f_{NL} to be momentum-independent. We have considered two typical ways to fix the initial state. One is to fix the state at a given time, common to all modes. The other is to fix the state for each mode at a given physical momentum. The former and the latter may be relevant when there was a phase transition, and when discussing trans-Planckian effects, respectively. We have spelled out the conditions for both cases and presented simple examples in which a large, scale-invariant f_{NL} is realized.

Naturally it is of great interest to see if these simple examples can be actually realized in any specific models of inflation. Researches in this direction are left for future study.

Acknowledgements

JG is grateful to the Yukawa Institute for Theoretical Physics at Kyoto University for hospitality while this work was under progress. We are grateful to Xingang Chen and Takahiro Tanaka for valuable comments on the first version of the paper. JG acknowledges the Max-Planck-Gesellschaft, the Korea Ministry of Education, Science and Technology, Gyeongsangbuk-Do and Pohang City for the support of the Independent Junior Research Group at the Asia Pacific Center for Theoretical Physics. This work was supported in part by the JSPS Grant-in-Aid for Scientific Research (A) No. 21244033.

-
- [1] A. H. Guth, Phys. Rev. D **23**, 347 (1981); K. Sato, Mon. Not. Roy. Astron. Soc. **195**, 467 (1981); A. D. Linde, Phys. Lett. B **108**, 389 (1982); A. Albrecht and P. J. Steinhardt, Phys. Rev. Lett. **48**, 1220 (1982).
 - [2] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith and R. S. Hill *et al.*, arXiv:1212.5225 [astro-ph.CO].
 - [3] [Planck Collaboration], arXiv:astro-ph/0604069.
 - [4] For a recent collection of reviews, see e.g. Class. Quant. Grav. **27**, “Focus section on non-linear and non-Gaussian cosmological perturbations” (2010); Adv. Astron. **2010**, “Testing the Gaussianity and Statistical Isotropy of the Universe” (2010).
 - [5] E. Komatsu and D. N. Spergel, Phys. Rev. D **63**, 063002 (2001) [astro-ph/0005036].
 - [6] J. M. Maldacena, JHEP **0305**, 013 (2003) [astro-ph/0210603].
 - [7] P. Creminelli and M. Zaldarriaga, JCAP **0410**, 006 (2004) [astro-ph/0407059].
 - [8] A. Achucarro, J. -O. Gong, G. A. Palma and S. P. Patil, arXiv:1211.5619 [astro-ph.CO].
 - [9] J. Ganc and E. Komatsu, JCAP **1012**, 009 (2010) [arXiv:1006.5457 [astro-ph.CO]]; S. Renaux-Petel, JCAP **1010**, 020 (2010) [arXiv:1008.0260 [astro-ph.CO]].
 - [10] M. H. Namjoo, H. Firouzjahi, M. Sasaki and , Europhys. Lett. **101**, 39001 (2013) [arXiv:1210.3692 [astro-ph.CO]]; X. Chen, H. Firouzjahi, M. H. Namjoo and M. Sasaki, arXiv:1301.5699 [hep-th].
 - [11] X. Chen, M. -x. Huang, S. Kachru and G. Shiu, JCAP **0701**, 002 (2007) [hep-th/0605045]; R. Holman and A. J. Tolley, JCAP **0805**, 001 (2008) [arXiv:0710.1302 [hep-th]].
 - [12] I. Agullo and L. Parker, Phys. Rev. D **83**, 063526 (2011) [arXiv:1010.5766 [astro-ph.CO]]; J. Ganc, Phys. Rev. D **84**, 063514 (2011) [arXiv:1104.0244 [astro-ph.CO]]; D. Chialva, JCAP **1210**, 037 (2012) [arXiv:1108.4203 [astro-ph.CO]]; N. Agarwal, R. Holman, A. J. Tolley and J. Lin, arXiv:1212.1172 [hep-th]. See also S. Kundu, JCAP **1202**, 005 (2012) [arXiv:1110.4688 [astro-ph.CO]].
 - [13] F. Arroja, A. E. Romano and M. Sasaki, Phys. Rev. D **84**, 123503 (2011) [arXiv:1106.5384 [astro-ph.CO]]; F. Arroja and M. Sasaki, JCAP **1208**, 012 (2012) [arXiv:1204.6489 [astro-ph.CO]].

- [14] T. Tanaka and Y. Urakawa, JCAP **1105**, 014 (2011) [arXiv:1103.1251 [astro-ph.CO]]; T. Tanaka and Y. Urakawa, arXiv:1209.1914 [hep-th].
- [15] J. Garriga and V. F. Mukhanov, Phys. Lett. B **458**, 219 (1999) [hep-th/9904176].
- [16] M. Sasaki, Prog. Theor. Phys. **76**, 1036 (1986).
- [17] C. T. Byrnes and J. -O. Gong, Phys. Lett. B **718**, 718 (2013) [arXiv:1210.1851 [astro-ph.CO]].
- [18] A. Vilenkin and L. H. Ford, Phys. Rev. D **26**, 1231 (1982).
- [19] See e.g. G. Shiu, J. Phys. Conf. Ser. **18**, 188 (2005) and references therein.